arXiv:1506.00255v1 [cs.DM] 31 May 2015

Critical and Maximum Independent Sets of a Graph

Adi Jarden
Department of Mathematics and Computer Science
Ariel University, Israel
jardena@ariel.ac.il

Vadim E. Levit
Department of Mathematics and Computer Science
Ariel University, Israel
levitv@ariel.ac.il

Eugen Mandrescu
Department of Computer Science
Holon Institute of Technology, Israel
eugen_m@hit.ac.il

Abstract

Let G be a simple graph with vertex set $V(G)$. A set $S \subseteq V(G)$ is *independent* if no two vertices from S are adjacent. By $\mathrm{Ind}(G)$ we mean the family of all independent sets of G, while $\mathrm{core}(G)$ and $\mathrm{corona}(G)$ denote the intersection and the union of all maximum independent sets, respectively.

The number $d(X) = |X| - |N(X)|$ is the *difference* of $X \subseteq V(G)$, and a set $A \in \mathrm{Ind}(G)$ is *critical* if $d(A) = \max\{d(I) : I \in \mathrm{Ind}(G)\}$ [23].

Let $\ker(G)$ and $\mathrm{diadem}(G)$ be the intersection and union, respectively, of all critical independent sets of G [13].

In this paper, we present various connections between critical unions and intersections of maximum independent sets of a graph. These relations give birth to new characterizations of König-Egerváry graphs, some of them involving $\ker(G)$, $\mathrm{core}(G)$, $\mathrm{corona}(G)$, and $\mathrm{diadem}(G)$.

Keywords: maximum independent set, maximum critical set, ker, core, corona, diadem, maximum matching, König-Egerváry graph.

1 Introduction

Throughout this paper G is a finite simple graph with vertex set $V(G)$ and edge set $E(G)$. If $X \subseteq V(G)$, then $G[X]$ is the subgraph of G induced by X. By $G - W$ we mean either the subgraph $G[V(G) - W]$, if $W \subseteq V(G)$, or the subgraph obtained by deleting the edge set W, for $W \subseteq E(G)$. In either case, we use $G - w$, whenever $W = \{w\}$. If $A, B \subseteq V(G)$, then (A, B) stands for the set $\{ab : a \in A, b \in B, ab \in E(G)\}$.

The *neighborhood* $N(v)$ of $v \in V(G)$ is the set $\{w : w \in V(G) \text{ and } vw \in E(G)\}$. In order to avoid ambiguity, we use also $N_G(v)$ instead of $N(v)$. The *neighborhood* $N(A)$ of $A \subseteq V(G)$ is $\{v \in V(G) : N(v) \cap A \neq \emptyset\}$, and $N[A] = N(A) \cup A$.

A set $S \subseteq V(G)$ is *independent* if no two vertices from S are adjacent, and by $\mathrm{Ind}(G)$ we mean the family of all the independent sets of G. An independent set of maximum size is a *maximum independent set* of G, and $\alpha(G) = \max\{|S| : S \in \mathrm{Ind}(G)\}$.

Theorem 1.1 *[1], [2] An independent set X is maximum if and only if every independent set S disjoint from X can be matched into X.*

For a graph G, let $\Omega(G)$ denote the family of all its maximum independent sets, $\mathrm{core}(G) = \bigcap\{S : S \in \Omega(G)\}$ [11], and $\mathrm{corona}(G) = \bigcup\{S : S \in \Omega(G)\}$ [3].

It is clear that $N(\mathrm{core}(G)) \subseteq V(G) - \mathrm{corona}(G)$, and there exist graphs satisfying $N(\mathrm{core}(G)) \neq V(G) - \mathrm{corona}(G)$ (for some examples, see the graphs from Figure 1, where $\mathrm{core}(G_1) = \{a, b\}$ and $\mathrm{core}(G_2) = \{x, y, z\}$).

The problem of whether $\mathrm{core}(G) \neq \emptyset$ is **NP**-hard [3].



Figure 1: $V(G_1) = \mathrm{corona}(G_1) \cup N(\mathrm{core}(G_1)) \cup \{d\}$, $V(G_2) = \mathrm{corona}(G_2) \cup N(\mathrm{core}(G_2))$.

A *matching* is a set M of pairwise non-incident edges of G. If $A \subseteq V(G)$, then $M(A)$ is the set of all the vertices matched by M with vertices belonging to A. A matching of maximum cardinality, denoted $\mu(G)$, is a *maximum matching.*

Lemma 1.2 (Matching Lemma) *[19] If $A \in \mathrm{Ind}(G), \Lambda \subseteq \Omega(G)$, and $|\Lambda| \geq 1$, then there exists a matching from $A - \bigcap\Lambda$ into $\bigcup\Lambda - A$.*

For $X \subseteq V(G)$, the number $|X| - |N(X)|$ is the *difference* of X, denoted $d(X)$. The *critical difference* $d(G)$ is $\max\{d(X) : X \subseteq V(G)\}$. The number $\max\{d(I) : I \in \mathrm{Ind}(G)\}$ is the *critical independence difference* of G, denoted $id(G)$. Clearly, $d(G) \geq id(G)$. It was shown in [23] that $d(G) = id(G)$ holds for every graph G. If A is an independent set in G with $d(X) = id(G)$, then A is a *critical independent set* [23].

For example, consider the graph G of Figure 2, where $X = \{v_1, v_2, v_3, v_4\}$ is a critical set, while $I = \{v_1, v_2, v_3, v_6, v_7\}$ is a critical independent set. Other critical sets are $\{v_1, v_2\}$, $\{v_1, v_2, v_3\}$, $\{v_1, v_2, v_3, v_4, v_6, v_7\}$.

It is known that finding a maximum independent set is an **NP**-hard problem [6]. Zhang proved that a critical independent set can be found in polynomial time [23].

Theorem 1.3 *[4] Each critical independent set is included in a maximum independent set.*



Figure 2: $\text{core}(G) = \{v_1, v_2, v_6, v_{10}\}$ is a critical set, since $d(\text{core}(G)) = 1 = d(G)$.

Theorem 1.3 leads to an efficient way of approximating $\alpha(G)$ [22]. Moreover, every critical independent set is contained in a maximum critical independent set, and such a maximum critical independent set can be found in polynomial time [9].

Theorem 1.4 *[9] There is a matching from $N(S)$ into S for every critical independent set S.*

It is well-known that $\alpha(G) + \mu(G) \leq |V(G)|$ holds for every graph G. Recall that if $\alpha(G) + \mu(G) = |V(G)|$, then G is a *König-Egerváry graph* [5, 21]. For example, each bipartite graph is a König-Egerváry graph as well. Various properties of König-Egerváry graphs can be found in [8, 12, 18]. It turns out that König-Egerváry graphs are exactly the graphs having a critical maximum independent set [10]. In [14] it was shown the following.

Lemma 1.5 *[14] $d(G) = \alpha(G) - \mu(G)$ holds for each König-Egerváry graph G.*

Using this finding, we have strengthened the characterization from [10].

Theorem 1.6 *[14] For a graph G, the following assertions are equivalent:*
(i) *G is a König-Egerváry graph;*
(ii) *there exists some maximum independent set which is critical;*
(iii) *each of its maximum independent sets is critical.*

For a graph G, let $\ker(G)$ be the intersection of all its critical independent sets [13], and $\text{diadem}(G) = \bigcup \{S : S \text{ is a critical independent set}\}$.

In this paper we present several properties of critical unions and intersections of maximum independent sets leading to new characterizations of König-Egerváry graphs, in terms of $\text{core}(G)$, $\text{corona}(G)$, and $\text{diadem}(G)$.

2 Preliminaries

Let G be the graph from Figure 2; the sets $X = \{v_1, v_2, v_3\}$, $Y = \{v_1, v_2, v_4\}$ are critical independent, and the sets $X \cap Y$, $X \cup Y$ are also critical, but only $X \cap Y$ is also independent. In addition, one can easily see that $\ker(G) = \{v_1, v_2\} \subseteq \text{core}(G)$, and $\ker(G)$ is a minimal critical independent set of G.

Theorem 2.1 *[13] For a graph G, the following assertions are true:*
(i) $\ker(G) \subseteq \text{core}(G)$;
(ii) *if A and B are critical in G, then $A \cup B$ and $A \cap B$ are critical as well;*
(iii) *G has a unique minimal independent critical set, namely,* $\ker(G)$.

Various properties of $\ker(G)$ and $\mathrm{core}(G)$ can be found in [15, 17, 20].

As an immediate consequence of Theorem 2.1, we have the following.

Corollary 2.2 *For every graph G, $\mathrm{diadem}(G)$ is a critical set.*

For instance, the graph G from Figure 2 has $\mathrm{diadem}(G) = \{v_1, v_2, v_3, v_4, v_6, v_7, v_{10}\}$, which is critical, but not independent.

The graph G_1 from Figure 1 has $d(G_1) = 1$ and $d(\mathrm{corona}(G_1)) = 0$, which means that $\mathrm{corona}(G_1)$ is not a critical set. Notice that G_1 is not a König-Egerváry graph. Combining Theorems 1.6 and 2.1*(ii)*, we deduce the following.

Corollary 2.3 *If G is a König-Egerváry graph, then both $\mathrm{core}(G)$ and $\mathrm{corona}(G)$ are critical sets. Moreover, $\mathrm{corona}(G) = \bigcup \{A : A$ is a maximum critical independent set$\}$.*

The converse of Corollary 2.3 is not necessarily true; e.g., the graph G_2 in Figure 1 is not a König-Egerváry graph, while $\mathrm{core}(G_2)$ and $\mathrm{corona}(G_2)$ are critical.

3 Unions and intersections of maximum independent sets

Theorem 3.1 *Let $\Lambda \subseteq \Omega(G)$, and $|\Lambda| \geq 1$. Then*

$$d\left(\bigcup \Lambda\right) = \left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| - |V(G)| \geq \max_{S \in \Lambda} d(S).$$

In particular,

$$d(\mathrm{corona}(G)) = |\mathrm{corona}(G)| + |\mathrm{core}(G)| - |V(G)| \geq 2\alpha(G) - |V(G)| = \max_{S \in \Omega(G)} d(S).$$

Proof. Every vertex in $\bigcup \Lambda - \bigcap \Lambda$ has a neighbor in $\bigcup \Lambda - \bigcap \Lambda$, since $\Lambda \subseteq \Omega(G)$. Therefore, $N\left(\bigcup \Lambda\right) = \left(\bigcup \Lambda - \bigcap \Lambda\right) \cup \left(V(G) - \bigcup \Lambda\right)$, which implies

$$\begin{aligned} d\left(\bigcup \Lambda\right) &= \left|\bigcup \Lambda\right| - \left|N\left(\bigcup \Lambda\right)\right| = \left|\bigcup \Lambda\right| - \left|\left(\bigcup \Lambda - \bigcap \Lambda\right) \cup \left(V(G) - \bigcup \Lambda\right)\right| = \\ &= \left|\bigcap \Lambda\right| - \left(|V(G)| - \left|\bigcup \Lambda\right|\right) = \left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| - |V(G)|. \end{aligned}$$

On the other hand, for every $S \in \Omega(G)$ we have

$$d(S) = \alpha(G) - (|V(G)| - \alpha(G)) = 2\alpha(G) - |V(G)|.$$

Since $\left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| \geq 2\alpha(G)$, we obtain

$$d\left(\bigcup \Lambda\right) = \left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| - |V(G)| \geq 2\alpha(G) - |V(G)| = d(S),$$

as required.

In particular, if $\Lambda = \Omega(G)$, then $\bigcup \Lambda = \text{corona}(G)$, $\bigcap \Lambda = \text{core}(G)$, and the conclusion follows. ■

Notice that if A is a critical independent set in a graph G having $d(G) > 0$, then $A \cap S \neq \emptyset$ holds for every $S \in \Omega(G)$, because $\emptyset \neq \ker(G) \subseteq A \cap \text{core}(G) \subseteq A \cap S$, according to Theorem 2.1*(i)*.

Proposition 3.2 *Let A be a critical independent set of a graph G with* $\ker(G) = \emptyset$*, and* $\Lambda = \{S \in \Omega(G) : A \cap S = \emptyset\}$*. Then* $\left|\bigcap \Lambda\right| \geq |A|$.

Proof. Let $S \in \Lambda$. Since A is critical and $d(G) = 0$, it follows that $|A| = |N(A)|$. By Theorem 1.1, there is a matching from A into S, because A is independent and disjoint from S. Consequently, we infer that $N(A) \subset S$. Hence, we obtain $\left|\bigcap \Lambda\right| \geq |N(A)| = |A|$, as required. ■

Theorem 3.3 *Let $\Lambda \subseteq \Omega(G)$, and $|\Lambda| \geq 1$.*

(i) *If $\bigcup \Lambda$ is critical, then $\left|N\left(\bigcap \Lambda\right)\right| + \left|\bigcup \Lambda\right| = |V(G)|$, and $\bigcap \Lambda$ is critical.*

(ii) *If $\bigcap \Lambda$ is critical, then*

$$\left|N\left(\bigcap \Lambda\right)\right| + \left|\bigcup \Lambda\right| \leq |V(G)|, \text{ and } d\left(\bigcap \Lambda\right) \geq 2\alpha(G) - |V(G)|.$$

Proof. *(i)* By definition of $d(G)$ and Theorem 3.1, we get

$$d(G) = d\left(\bigcup \Lambda\right) = \left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| - |V(G)| \geq d\left(\bigcap \Lambda\right) = \left|\bigcap \Lambda\right| - \left|N\left(\bigcap \Lambda\right)\right|.$$

Hence we infer that $\left|N\left(\bigcap \Lambda\right)\right| + \left|\bigcup \Lambda\right| \geq |V(G)|$. Thus, $\left|N\left(\bigcap \Lambda\right)\right| + \left|\bigcup \Lambda\right| = |V(G)|$, because $\left(N\left(\bigcap \Lambda\right)\right) \cap \left(\bigcup \Lambda\right) = \emptyset$.

Moreover, we deduce that

$$d\left(\bigcup \Lambda\right) = \left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| - |V(G)| = \left|\bigcap \Lambda\right| - \left|N\left(\bigcap \Lambda\right)\right| = d\left(\bigcap \Lambda\right),$$

i.e., $\bigcap \Lambda$ is a critical set.

(ii) By definition of $d(G)$ and Theorem 3.1, we have

$$d(G) = d\left(\bigcap \Lambda\right) = \left|\bigcap \Lambda\right| - \left|N\left(\bigcap \Lambda\right)\right| \geq$$
$$\geq d\left(\bigcup \Lambda\right) = \left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| - |V(G)| \geq 2\alpha(G) - |V(G)|,$$

which completes the proof. ■

In particular, taking $\Lambda = \Omega(G)$ in Theorem 3.3, we obtain the following.

Corollary 3.4 *If* $\text{corona}(G)$ *is a critical set, then* $|\text{corona}(G)| + |N(\text{core}(G))| = |V(G)|$ *and* $\text{core}(G)$ *is critical.*

Notice that if core(G) is critical, then corona(G) is not necessarily critical. For example, the graph G_1 from Figure 1 has $d(G_1) = d(\text{core}(G_1)) = 1$, while corona$(G_1)$ is not a critical set.

Theorem 3.5 *If G is a König-Egerváry graph, then*

(i) *[7]* $\left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| = 2\alpha(G)$ *holds for every family* $\Lambda \subseteq \Omega(G)$, $|\Lambda| \geq 1$*;*

(ii) *[19]* $|\text{corona}(G)| + |\text{core}(G)| = 2\alpha(G)$.

Proof. *(i)* By Theorems 1.6 and 2.1*(ii)*, both $\bigcup \Lambda$ and $\bigcap \Lambda$ are critical sets. According to Lemma 1.5, we have

$$d\left(\bigcup \Lambda\right) = \left|\bigcup \Lambda\right| - \left|N\left(\bigcup \Lambda\right)\right| = \alpha(G) - \mu(G),$$
$$d\left(\bigcap \Lambda\right) = \left|\bigcap \Lambda\right| - \left|N\left(\bigcap \Lambda\right)\right| = \alpha(G) - \mu(G).$$

Hence, $\left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| = 2\alpha(G) - 2\mu(G) + \left|N\left(\bigcup \Lambda\right)\right| + \left|N\left(\bigcap \Lambda\right)\right|$.

By Theorem 3.3*(i)*, we infer that

$$\left|N\left(\bigcup \Lambda\right)\right| + \left|N\left(\bigcap \Lambda\right)\right| = \left|N\left(\bigcup \Lambda\right)\right| + |V(G)| - \left|\bigcup \Lambda\right| =$$
$$= \left|N\left(\bigcup \Lambda\right)\right| + \alpha(G) + \mu(G) - \left|\bigcup \Lambda\right| =$$
$$= \alpha(G) + \mu(G) - d\left(\bigcup \Lambda\right) = 2\mu(G).$$

Consequently, we obtain $\left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| = 2\alpha(G)$, as claimed.

(ii) It follows from Part *(i)*, by taking $\Lambda \subseteq \Omega(G)$. ■

The graph G_2 from Figure 1 has $|\text{corona}(G_2)| + |\text{core}(G_2)| = 13 > 12 = 2\alpha(G_2)$. On the other hand, there is a non-König-Egerváry graph, namely G_1 in Figure 1, that satisfies $|\text{corona}(G_1)| + |\text{core}(G_1)| = 10 = 2\alpha(G_1)$.

If $\bigcap \Lambda$ is a critical set, then $\bigcup \Lambda$ is not necessarily critical. For instance, consider the graph G from Figure 3, and $\Lambda = \{S_1, S_2\}$, where $S_1 = \{x, y, u\}$ and $S_2 = \{x, y, w\}$. Clearly, $\bigcap \Lambda = \{x, y\} = \text{core}(G)$ is critical, while $\bigcup \Lambda = \{x, y, u, w\}$ is not a critical set.



Figure 3: core$(G) = \ker(G) = \{x, y\}$.

Theorem 3.6 *Let $\Lambda \subseteq \Omega(G)$, and $|\Lambda| \geq 1$. Then G is a König-Egerváry graph if and only if $\bigcup \Lambda$ is critical and $\left|\bigcap \Lambda\right| + \left|\bigcup \Lambda\right| = 2\alpha(G)$.*

Proof. Combining Theorems 1.6 and 2.1*(ii)*, we infer that $\bigcup\Lambda$ is critical. The equality $\left|\bigcap\Lambda\right| + \left|\bigcup\Lambda\right| = 2\alpha(G)$ holds by Theorem 3.5*(i)*.

Conversely, according to Theorem 3.3*(i)*, the set $\bigcap\Lambda$ is critical. Hence, by Theorem 1.4, there exists a matching from $N\left(\bigcap\Lambda\right)$ into $\bigcap\Lambda$. Theorem 3.3*(i)* ensures that $\left|N\left(\bigcap\Lambda\right)\right| + \left|\bigcup\Lambda\right| = |V(G)|$, which means that $\bigcup\Lambda \cup N\left(\bigcap\Lambda\right) = V(G)$. To complete the proof that G is a König-Egerváry graph, one has to find a matching from $\bigcup\Lambda - S$ into $S - \bigcap\Lambda$ for some maximum independent set $S \in \Lambda$. Actually, in accordance with Lemma 1.2, there is a matching, say M, from $S - \bigcap\Lambda$ to $\bigcup\Lambda - S$. Since $\left|\bigcap\Lambda\right| + \left|\bigcup\Lambda\right| = 2\alpha(G)$ and $|S| = \alpha(G)$, we infer that $\left|S - \bigcap\Lambda\right| = \left|\bigcup\Lambda - S\right|$. Consequently, M is a perfect matching, and this shows that M is also a matching from $\bigcup\Lambda - S$ into $S - \bigcap\Lambda$, as required. ■

Remark 3.7 *If* $\bigcap\Lambda$ *is critical and* $\left|\bigcap\Lambda\right| + \left|\bigcup\Lambda\right| = 2\alpha(G)$*, then* G *is not necessarily a König-Egerváry graph. For example, let* G *be the graph from Figure 3, and* $\Lambda = \{S_1, S_2\}$*, where* $S_1 = \{x, y, u\}, S_2 = \{x, y, w\}$*. Hence,* $\bigcap\Lambda = \{x, y\} = \ker(G)$ *is a critical set,* $\left|\bigcap\Lambda\right| + \left|\bigcup\Lambda\right| = 6 = 2\alpha(G)$*, while* G *is not a König-Egerváry graph. Clearly, the set* $\bigcup\Lambda = \{x, y, u, w\}$ *is not critical.*



Figure 4: G is a non-König-Egerváry graph with $\text{core}(G) = \ker(G) = \{x, y\}$.

If A_1, A_2 are independent sets such that $A_1 \cup A_2$ is critical, then A_1 and A_2 are not necessarily critical. For instance, consider the graph G from Figure 4, where $A_1 \cup A_2 = \{u, v, x, y\}$ is a critical set, while none of $A_1 = \{u, x\}$ and $A_2 = \{v, y\}$ is critical. The case is different whenever the two independent sets are also maximum.

Corollary 3.8 *The following assertions are equivalent:*

(i) *G is a König-Egerváry graph;*

(ii) *for every $S_1, S_2 \in \Omega(G)$, the set $S_1 \cup S_2$ is critical;*

(iii) *there exist $S_1, S_2 \in \Omega(G)$, such that $S_1 \cup S_2$ is critical.*

Proof. *(i)* ⇒ *(ii)* It follows combining Theorem 1.6*(iii)* and Theorem 2.1*(ii)*.

(ii) ⇒ *(iii)* Clear.

(iii) ⇒ *(i)* It is true according to Theorem 3.6, because

$$\left|\bigcap\Lambda\right| + \left|\bigcup\Lambda\right| = |S_1 \cup S_2| + |S_1 \cap S_2| = |S_1| + |S_2| = 2\alpha(G)$$

is automatically valid for every family $\Lambda \subseteq \Omega(G)$ with $|\Lambda| = 2$. ■

Remark 3.9 *If G is a König-Egerváry graph, then $S \cup A$ is not necessarily critical for every $S \in \Omega(G)$ and $A \in \mathrm{Ind}(G)$. For instance, consider the graph G in Figure 5, and $S = \{a, b, c, d\} \in \Omega(G)$. The sets $A_1 = \{v\}$ and $A_2 = \{w\}$ are independent, $S \cup A_1$ is critical (because $N(S \cup A_1) = \{u, v, w, a\}$), while $S \cup A_2$ is not critical (as $N(S \cup A_2) = \{u, v, w, c, d\}$).*



Figure 5: G is a König-Egerváry graph with $d(G) = 1$.

Corollary 3.10 *The following assertions are equivalent:*

(i) *G is a König-Egerváry graph;*

(ii) *for every $S \in \Omega(G)$ there exists $A \in \mathrm{Ind}(G)$, such that the set $S \cup A$ is critical;*

(iii) *there are $S \in \Omega(G)$ and $A \in \mathrm{Ind}(G)$, such that the set $S \cup A$ is critical.*

Proof. *(i) ⇒ (ii)* By Theorem 1.6*(iii)*, we know that every $S \in \Omega(G)$ is critical. Hence, $S \cup A$ is critical for any $A \subseteq S$.

(ii) ⇒ (iii) Clear.

(iii) ⇒ (i) If $A \subseteq S$, the result follows by Theorem 1.6. Otherwise, we can suppose that $S \cap A = \emptyset$. By Theorem 1.1, we know that $|N(A) \cap S| \geq |A|$. Since

$$|N(S \cup A)| \geq |N(A) \cap S| + |N(S)| \geq |A| + |N(S)|,$$

we obtain

$$\begin{aligned} d(G) = d(S \cup A) &= |S \cup A| - |N(S \cup A)| \\ &\leq (|S| + |A|) - (|A| + |N(S)|) = d(S). \end{aligned}$$

Therefore, $d(S) = d(G)$, i.e., S is a critical set. According to Theorem 1.6, G is a König-Egerváry graph. ■

4 $\ker(G)$ and $\mathrm{diadem}(G)$ in König-Egerváry graphs

Theorem 4.1 *If G is a König-Egerváry graph, then*

(i) $\mathrm{diadem}(G) = \mathrm{corona}(G)$, *while* $\mathrm{diadem}(G) \subseteq \mathrm{corona}(G)$ *is true for every graph;*

(ii) $|\ker(G)| + |\mathrm{diadem}(G)| \leq 2\alpha(G)$.

Proof. *(i)* Every $S \in \Omega(G)$ is a critical set, by Theorem 1.6. Hence we deduce that $\mathrm{corona}(G) \subseteq \mathrm{diadem}(G)$. On the other hand, for every graph each critical independent set is included in a maximum independent set, according to Theorem 1.3. Thus, we infer that $\mathrm{diadem}(G) \subseteq \mathrm{corona}(G)$. Consequently, the equality $\mathrm{diadem}(G) = \mathrm{corona}(G)$ holds.

(ii) It follows by combining Part *(i)*, Theorem 3.5*(ii)* and Theorem 2.1*(i)*. ■



Figure 6: G_1 and G_2 are König-Egerváry graphs. $ker(G_1) = \{x, y\}$ and $ker(G_2) = \emptyset$.

The König-Egerváry graphs from Figure 6 satisfy $|\ker(G)| + |\text{diadem}(G)| < 2\alpha(G)$.

The graph G_1 from Figure 7 is a non-bipartite König-Egerváry graph, such that $\ker(G_1) = \text{core}(G_1)$ and $\text{diadem}(G_1) = \text{corona}(G_1)$. The combination of $\text{diadem}(G) \subsetneqq \text{corona}(G)$ and $\ker(G) = \text{core}(G)$ is realized by the non-König-Egerváry graph G_2 from Figure 7, because $\ker(G_2) = \text{core}(G_2)$ and $\text{diadem}(G_2) \cup \{z, t, v, w\} = \text{corona}(G_2)$.



Figure 7: $\text{core}(G_1) = \{a, b\}$ and $\text{core}(G_2) = \{x, y\}$.

The graph G_2 from Figure 1 has : $\text{corona}(G_2) = V(G_2) - \{v_1, v_6\}$ and $N(\text{corona}(G_2)) = V(G_2) - \{x, y, z\}$. Hence, $d(\text{corona}(G_2)) = 1 = d(G_2)$, i.e., $\text{corona}(G_2)$ is a critical set, while $\text{diadem}(G_2) = \{x, y, z, v_2, v_3, v_5\} \subsetneqq \text{corona}(G_2)$. Thus, the graph G_2 from Figure 1 shows that it is possible for a graph to have $\text{diadem}(G) \subsetneqq \text{corona}(G)$ and $\ker(G) \subsetneqq \text{core}(G)$. On the other hand, the graph G_2 from Figure 1 gives an example where not every critical set is a subset of $\text{diadem}(G)$.

Corollary 4.2 *The following assertions are equivalent:*
(i) *G is a König-Egerváry graph;*
(ii) $\text{diadem}(G) = \text{corona}(G)$ *and* $|\text{core}(G)| + |\text{corona}(G)| = 2\alpha(G)$*;*
(iii) $\text{corona}(G)$ *is a critical set and* $|\text{core}(G)| + |\text{corona}(G)| = 2\alpha(G)$.

Proof. *(i) ⇒ (ii)* It is true, by applying Theorem 4.1*(i)* and Theorem 3.5*(ii)*.
(ii) ⇒ (iii) It follows from Corollary 2.2.
(iii) ⇒ (i) Take $\Lambda = \Omega(G)$ and use Theorem 3.6. ■

Notice that the graph G_1 from Figure 1 satisfies $|\text{core}(G_1)| + |\text{corona}(G_1)| = 2\alpha(G_1)$, while $d(\text{corona}(G_1)) = 0 < d(G_1) = 1$, i.e., $\text{corona}(G_1)$ is not a critical set, because $\text{corona}(G_1) = V(G_1) - \{c, d\}$ and $N(\text{corona}(G_1)) = V(G_1) - \{a, b\}$. On the other hand, the graph G_2 from Figure 1 satisfies $|\text{core}(G_2)| + |\text{corona}(G_2)| = 13 > 12 = 2\alpha(G_2)$, while $\text{corona}(G_2)$ is a critical set.

5 Conclusions

In this paper we focus on interconnections between critical unions and intersections of maximum independent sets, with emphasis on König-Egerváry graphs. In [19] we showed that $2\alpha(G) \leq |\text{core}(G)| + |\text{corona}(G)|$ is true for every graph, while the equality $\text{diadem}(G) = \text{corona}(G)$ holds for each König-Egerváry graph G, by Theorem 4.1*(i)*.

According to Theorem 2.1*(i)*, $\ker(G) \subseteq \mathrm{core}(G)$ for every graph. On the other hand, Theorem 1.3 implies the inclusion $\mathrm{diadem}(G) \subseteq \mathrm{corona}(G)$. Hence

$$|\ker(G)| + |\mathrm{diadem}(G)| \leq |\mathrm{core}(G)| + |\mathrm{corona}(G)|$$

for each graph G. These remarks together with Theorem 4.1*(ii)* motivate the following.

Conjecture 5.1 *$|\ker(G)| + |diadem(G)| \leq 2\alpha(G)$ is true for every graph G.*

When it is proved one can conclude that the following inequalities:

$$|\ker(G)| + |\mathrm{diadem}(G)| \leq 2\alpha(G) \leq |\mathrm{core}(G)| + |\mathrm{corona}(G)|$$

hold for every graph G.

Theorem 4.1 claims that $\mathrm{diadem}(G) = \mathrm{corona}(G)$ is a necessary condition for G to be a König-Egerváry graph, while Corollary 4.2 shows that, apparently, this equality is not enough. These facts motivate the following.

Conjecture 5.2 *If* $\mathrm{diadem}(G) = \mathrm{corona}(G)$, *then G is a König-Egerváry graph.*

The graphs in Figure 8 are non-König-Egerváry graphs; $\mathrm{core}(G_1) = \{a, b, c, d\}$ and it is a critical set, while $\mathrm{core}(G_2) = \{x, y, z, w\}$ and it is not critical.



Figure 8: Both G_1 and G_2 are non-König-Egerváry graphs.

By Corollary 2.3, $\mathrm{core}(G)$ is a critical set for every König-Egerváry graph. It justifies the following.

Problem 5.3 *Characterize graphs such that* $\mathrm{core}(G)$ *is a critical set.*

It is known that the sets $\ker(G)$ and $\mathrm{core}(G)$ coincide for bipartite graphs [16]. Notice that there are non-bipartite graphs enjoying the equality $\ker(G) = \mathrm{core}(G)$; e.g., the graphs from Figure 9, where only G_1 is a König-Egerváry graph.



Figure 9: $\mathrm{core}(G_1) = \ker(G_1) = \{x, y\}$ and $\mathrm{core}(G_2) = \ker(G_2) = \{a, b\}$.

There is a non-bipartite König-Egerváry graph G, such that $\ker(G) \neq \mathrm{core}(G)$. For instance, the graph G_1 from Figure 6 has $\ker(G_1) = \{x, y\}$, while $\mathrm{core}(G_1) = \{x, y, u, v\}$. The graph G_2 from Figure 6 has $\ker(G_2) = \emptyset$, while $\mathrm{core}(G_2) = \{w\}$. We propose the following.

Problem 5.4 *Characterize (König-Egerváry) graphs satisfying* $\ker(G) = \mathrm{core}(G)$.

References

[1] C. Berge, *Some common properties for regularizable graphs, edge-critical graphs and B-graphs*, Lecture Notes in Computer Science **108** (1981) 108–123.

[2] C. Berge, *Graphs*, North-Holland, New York, 1985.

[3] E. Boros, M. C. Golumbic, V. E. Levit, *On the number of vertices belonging to all maximum stable sets of a graph*, Discrete Applied Mathematics **124** (2002) 17–25.

[4] S. Butenko, S. Trukhanov, *Using critical sets to solve the maximum independent set problem*, Operations Research Letters **35** (2007) 519–524.

[5] R. W. Deming, *Independence numbers of graphs - an extension of the König-Egerváry theorem*, Discrete Mathematics **27** (1979) 23–33.

[6] M. Garey, D. Johnson, *Computers and intractability*, W. H. Freeman and Company, New York, 1979.

[7] A. Jarden, V. E. Levit, E. Mandrescu, *Monotonic properties of collections of maximum independent sets of a graph*, arXive:1268080 [cs.DM] (2015) 15pp.

[8] E. Korach, T. Nguyen, B. Peis, *Subgraph characterization of red/blue-split graphs and König-Egerváry graphs*, in: Proceedings of the Seventeenth Annual ACM–SIAM Symposium on Discrete Algorithms, ACM Press, 2006, 842–850.

[9] C. E. Larson, *A note on critical independence reductions*, Bulletin of the Institute of Combinatorics and its Applications **5** (2007) 34–46.

[10] C. E. Larson, *The critical independence number and an independence decomposition*, European Journal of Combinatorics **32** (2011) 294–300.

[11] V. E. Levit, E. Mandrescu, *Combinatorial properties of the family of maximum stable sets of a graph*, Discrete Applied Mathematics **117** (2002) 149–161.

[12] V. E. Levit, E. Mandrescu, *On α^{+}-stable König-Egerváry graphs*, Discrete Mathematics **263** (2003) 179–190.

[13] V. E. Levit, E. Mandrescu, *Vertices belonging to all critical independent sets of a graph*, SIAM Journal on Discrete Mathematics **26** (2012) 399–403.

[14] V. E. Levit, E. Mandrescu, *Critical independent sets and König-Egerváry graphs*, Graphs and Combinatorics **28** (2012) 243–250.

[15] V. E. Levit, E. Mandrescu, *On the core of a unicyclic graph*, Ars Mathematica Contemporanea **5** (2012) 325–331.

[16] V. E. Levit, E. Mandrescu, *Critical sets in bipartite graphs*, Annals of Combinatorics **17** (2013) 543–548.

[17] V. E. Levit, E. Mandrescu, *On the structure of the minimum critical independent set of a graph*, Discrete Mathematics **313** (2013) 605–610.

[18] V. E. Levit, E. Mandrescu, *On maximum matchings in König-Egerváry graphs*, Discrete Applied Mathematics **161** (2013) 1635–1638.

[19] V. E. Levit, E. Mandrescu, *A set and collection lemma*, The Electronic Journal of Combinatorics **21** (2014) #P1.40.

[20] V. E. Levit, E. Mandrescu, *On the intersection of all critical sets of a unicyclic graph*, Discrete Applied Mathematics **162** (2014) 409–414.

[21] F. Sterboul, *A characterization of the graphs in which the transversal number equals the matching number*, Journal of Combinatorial Theory B **27** (1979) 228–229.

[22] S. Trukhanov, *Novel approaches for solving large-scale optimization problems on graphs*, Ph.D. Thesis, University of Texas, 2008.

http://repository.tamu.edu/bitstream/handle/1969.1/ETD-TAMU-2986/TRUKHANOV-DISSERTATIO

[23] C. Q. Zhang, *Finding critical independent sets and critical vertex subsets are polynomial problems*, SIAM Journal on Discrete Mathematics **3** (1990) 431–438.